

# Journal of Nephrothology



## Extensive renal cortical necrosis in a postpartum patient after COVID-19 infection

Ata Abbasi<sup>1,2\*</sup>, Khadijeh Makhdoomi<sup>3</sup>, Shadi Babaei<sup>2</sup>, Behzad Banaie<sup>2</sup>, Leila Mahmoudzadeh<sup>2</sup>, Sepideh Rahimi<sup>4</sup>, Parham Porteghali<sup>5</sup>

<sup>1</sup>Cellular and Molecular Research Center, Cellular and Molecular Medicine Institute, Urmia University of Medical Sciences, Urmia, Iran

<sup>2</sup>Department of Pathology, Urmia University of Medical Sciences, Urmia, Iran

<sup>3</sup>Nephrology and Kidney Transplant Research Center, Clinical Research Institute, Urmia University of Medical Sciences, Urmia, Iran

<sup>4</sup>Department of Pathology, Guilan University of Medical Sciences, Rasht, Iran

<sup>5</sup>Department of Internal Medicine, Guilan University of Medical Sciences, Rasht, Iran

### ARTICLE INFO

*Article type:*  
Case Report

*Article history:*  
Received: 28 May 2022  
Accepted: 14 April 2023  
Published online: 9 May 2023

*Keywords:*  
COVID-19, Acute kidney injury,  
Renal cortical necrosis, Pregnancy

### ABSTRACT

It has been more than two years since the COVID-19 pandemic affects the world. COVID-19 affects other organs in addition to the respiratory system. Renal cortical necrosis is a severe kidney injury due to decreased renal arterial perfusion induced by vascular spasm, microvascular injury and endovascular thrombosis. Acute kidney injury is a common finding in COVID-19; however, renal cortical necrosis is rare. To the best of our knowledge, only one case of renal cortical necrosis has been reported in the literature.

### *Implication for health policy/practice/research/medical education:*

COVID-19 infection is accompanied by several complications in addition to the respiratory disorders. Sufficient knowledge about all probable complications is needed for correct and on-time management of patients.

*Please cite this paper as:* Abbasi A, Makhdoomi K, Babaei S, Banaie B, Mahmoudzadeh L, Rahimi S, Porteghali P. Extensive renal cortical necrosis in a postpartum patient after COVID-19 infection. J Nephrothol. 2023;x(x):e18380. DOI: 10.34172/jnp.2023.18380.

### Introduction

It is more than two years that the world is affected by COVID-19 pandemic which has caused huge mortality and morbidity worldwide. Although its main manifestations are fever and respiratory distress; however, now it is clear that this virus can target almost all of the organs during the infection. Several complications have been reported for COVID-19 infection, some were very mild such as skin rashes but some were life threatening. Renal cortical necrosis is a fatal kind of kidney injury with serious outcomes which may lead to end-stage renal disease or death(1). Renal cortical necrosis accounts for 2% and 6-7% of all cases of acute renal failure in developed nations and developing countries, respectively (2). It is mostly bilateral and unilateral cortical necrosis is rare (3). Renal cortical necrosis is the result of decreased renal arterial

perfusion due to vascular spasm, microvascular injury and endovascular thrombosis (1). Histologically total ischemic necrosis of all of the renal cortical elements including glomeruli, blood vessels and tubules of the affected area is typical feature of Renal cortical necrosis (1). According to histologic findings two types of renal cortical necrosis have been described, diffuse and patchy cortical necrosis (1).

Among hospitalized patients with coronavirus infection, acute kidney injury (AKI) is a common finding (4); however, cortical necrosis is very rare and to the best of our knowledge, only one case of renal cortical necrosis has been reported in the literature (5). Here, we report the second case of renal cortical necrosis in a patient with COVID-19 infection.

\*Corresponding author: Ata Abbasi,  
Email: aabbasi@alumnum.tums.ac.ir

### Case Presentation

The patient was a previous healthy 37-years-old woman. She had an uncomplicated term vaginal delivery 12 days before her admission to our hospital. Three days after natural vaginal delivery, the patient returned to emergency with dyspnea and cyanosis. On initial physical examination she was found to be conscious and oriented with normal vital signs. Body temperature was 37.1, respiratory rate was 30 with O<sub>2</sub> saturation of 93%. According to positive polymerase chain reaction (PCR) test for COVID-19 and computerized tomography (CT) scan findings she was treated for COVID-19 infection. During admission, the patient's clinical condition worsened and laboratory findings showed uremia, elevated serum creatinine level, hyperphosphatemia and hypocalcemia (laboratory findings are mentioned in Table 1).

Considering the laboratory data advanced renal failure and rhabdomyolysis was suspected and she underwent hemodialysis through jugular vein. Ultrasonography of kidneys showed increased renal cortical echo pattern and more prominent corticomedullary differentiation with a simple cortical cyst measuring 21 mm in greatest diameter in the middle pole of the left kidney. Color Doppler

ultrasonography of the kidneys showed normal blood flow of renal arteries. Due to patient's progressive course renal biopsy was conducted for further evaluation of the kidney.

Renal biopsy showed extensive coagulative (ischemic) necrosis in about 60% of renal tissue mostly in cortical area involving both tubules and glomeruli. There was interstitial mixed inflammatory cells infiltration (about 60%) (mostly lymphocytes and rare neutrophils), hemosiderin deposition and mild edema in interstitium (Figures 1-3).


The patient was stabilized in hospital with corticosteroid therapy, supportive care and hemodialysis. After rigorous treatment efforts, the patient survived but the renal damage remained persistent. She was discharged from the hospital with serum creatinine level of 7.4 mg/dL with scheduled hemodialysis three times-per-week.

### Discussion


We report the case of previously a healthy 37-year-old female with history of normal vaginal delivery and COVID-19 infection with CT scan findings of pulmonary involvement and laboratory findings of uremia and hyperphosphatemia who underwent

**Table 1.** Laboratory finding of the patient


	Laboratory test	Value	Unit	Reference value
Hematology	Hemoglobin	10.8	g/dl	11-16
	Total leukocyte count	15.72	×1000/μL	4-10
	Platelet count	138	×1000/μL	140-440
	Uncorrected reticulocyte count	0.7	%	0.5-1.5
Biochemistry	Blood urea	204	mg/dL	0.6-1.4
	Uric acid	10.58	mg/dL	3.5-7.2
	Serum Creatinine	8	mg/dL	0.6-1.4
	Serum sodium	134	mEq/L	136-145
	Serum potassium	5.4	mEq/L	3.5-5.5
	Serum phosphorus	10.77	mg/dL	2.8-4.5
	Serum calcium	8	mg/dL	8.5-10.5
	Serum magnesium	2.3	mg/dL	1.8-2.6
	Serum total bilirubin	2.21	mg/dL	0.1-1.2
	Serum direct bilirubin	0.13	mg/dL	Less than 0.3
	Serum albumin	2.64	gr/dl	3.4-5.4
	Aspartate aminotransferase (AST)	15	U/L	Less than 37
	Alanine aminotransferase (ALT)	9	U/L	Less than 41
	Alkaline phosphatase (ALP)	193	U/L	80-306
Serum lactate dehydrogenase (LDH)	4055	U/L	140-280	
Urine	Urine protein	Trace		Negative
	Urine blood	Negative		Negative
	Red blood cell	3-4	Cell/hpf	Less than 3/hpf
	Urine 24 h protein	399	mg/day	Less than 150
	Urine 24 h volume	700	mL/day	800-1800
	Urine 24 h creatinine	175	mg/day	800-2000
Others	HBsAg	Negative		Negative
	ANA (antinuclear antibody)	0.2	Index	Less than 1.5
	CH50	38	U/mL	42-95



**Figure 1.** Renal biopsy with 3 totally necrotic glomeruli and chronic inflammatory cells infiltration of interstitium.



**Figure 2.** Extensive necrosis of glomeruli and tubules with interstitial chronic inflammatory cells infiltration.



**Figure 3.** Extensive necrosis of all elements in kidney nephrons.

hemodialysis with initial suspicion of rhabdomyolysis. Diagnosis of renal cortical necrosis was made after biopsy and histopathologic evaluation. Cortical necrosis is a rare cause of AKI. Renal cortical necrosis is mostly a result of decreased renal arterial perfusion (3); however, our patient had a normal arterial blood flow of both kidneys on color Doppler ultrasonography. Renal cortical necrosis of this patient can be caused due to microvascular injury of the kidney due to covid19 infection.

The most common reported causes of intensive care unit admission in COVID-19 patients were either hypoxemic respiratory failure needing mechanical ventilation or hypotension and data on AKI are little (6).

Several clinical and autopsy reports of COVID-19 patients from China and the United States revealed increased clotting and disseminated intravascular coagulation resulting in microangiopathy of other organ systems such as splenic infarction (6).

The incidence of AKI in COVID-19 patients is reported from 0.5 to 80 percent. The three most common histopathologic findings of kidney biopsies in COVID-19 patients were acute tubular injury, collapsing glomerulopathy and thrombotic microangiopathy and these data about pathology of AKI in COVID-19 patients are primarily derived from case reports and case series (7,8).

### Conclusion

The rarity of renal cortical necrosis, its important adverse outcomes and its unknown effects on vital systems prompts us to report this case. It is important to emphasize that early recognition and accurate management of COVID-19 infection is necessary to reduce its adverse complications.

### Authors' contribution

Conceptualization: AA, KM.

Methodology: AA.

Validation: AA, LM.

Investigation: SB, BB.

Resources: KM.

Data curation: AA, KM.

Writing—original draft preparation: SR, PP.

Writing—review and editing: AA, LM.

Visualization: AA, KM.

Supervision: AA, KM.

### Conflicts of interest

The authors declare that they have no competing interests.

### Ethical issues

This case report was conducted in accord with the World Medical Association Declaration of Helsinki. Written informed consent was obtained from the patient for publication of this report.

Ethical issues (including plagiarism, data fabrication, double publication) have been completely Observed by the authors.

### Funding/Support

No funds were received for this study.

### References

1. Prakash J, Singh VP. Changing picture of renal cortical necrosis in acute kidney injury in developing country. *World J Nephrol.* 2015;4):480-6. doi: 10.5527/wjn.v4.i5.480.

2. Prakash J, Vohra R, Wani IA, Murthy AS, Srivastva PK, Tripathi K, et al. Decreasing incidence of renal cortical necrosis in patients with acute renal failure in developing countries: a single-centre experience of 22 years from Eastern India. *Nephrol Dial Transplant*. 2007;22:1213-7. doi: 10.1093/ndt/gfl761.
3. Haaskjold YL, Nylund K, Skrunes R. A rare case of unilateral renal cortical necrosis. *Case Rep Nephrol Dial*. 2021;11:63-8. doi: 10.1159/000512228.
4. Ng JH, Bijol V, Sparks MA, Sise ME, Izzedine H, Jhaveri KD. Pathophysiology and pathology of acute kidney injury in patients with COVID-19. *Adv Chronic Kidney Dis*. 2020;27:365-76. doi: 10.1053/j.ackd.2020.09.003.
5. COVID-19 Autopsy. The first COVID-19 autopsy in Spain performed during the early stages of the pandemic. *Rev Esp Patol*. 2020;53:182-7. doi: 10.1016/j.patol.2020.05.004.
6. Battle D, Soler MJ, Sparks MA, Hiremath S, South AM, Welling PA, et al. Acute kidney injury in COVID-19: emerging evidence of a distinct pathophysiology. *J Am Soc Nephrol*. 2020;31:1380-3. doi: 10.1681/asn.2020040419.
7. Sharma P, Ng JH, Bijol V, Jhaveri KD, Wanchoo R. Pathology of COVID-19-associated acute kidney injury. *Clin Kidney J*. 2021;14:i30-i9. doi: 10.1093/ckj/sfab003.
8. de Oliveira P, Cunha K, Neves P, Muniz M, Gatto G, Salgado Filho N, et al. Renal morphology in coronavirus disease: a literature review. *Medicina (Kaunas)*. 2021;57(3):258. doi: 10.3390/medicina57030258.

**Copyright** © 2023 The Author(s); Published by Society of Diabetic Nephropathy Prevention. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.